

[psng]mize  
times  
[dummy 1893]dum et al. XMM-Newton  
document [The nature of X-ray selected EROs] The nature of X-ray selected EROs  
[J. A. Stevens et al.] J. A. Stevens<sup>1,2</sup>, M. J. Page<sup>2</sup>, R. J. Ivison<sup>1</sup>, Ian Smail<sup>3</sup>, I. Lehmann<sup>4,5</sup>, G.  
Hasinger<sup>4,5</sup> and G. Szokoly<sup>4,5</sup>  
draft 1.0

abstract We report on the X-ray, optical, near-infrared, submillimetre and radio properties of five Extremely Red Objects (EROs) selected at X-ray wavelengths by XMM-Newton in the Lockman Hole field. They all have enough counts in the X-ray band to allow spectral fitting: four are most probably obscured, Compton-thin AGN with redshift dependent absorbing column densities of  $10^{22} - 10^{24} \text{ cm}^{-2}$ , whilst the fifth is best fitted by a thermal spectrum and is likely to be a massive elliptical galaxy in a deep gravitational potential. Their optical/near-infrared colours and sizes suggest that X-ray selected EROs comprise a mixture of dusty ‘starburst’ galaxies and non-dusty galaxies that are dominated by either star-light or light from an active nucleus. The colour diagnostics are supported by the submillimetre and radio data: the two AGN with ‘starburst’ colours have submillimetre or radio flux densities that imply large star-formation rates, whilst those with ‘elliptical’ colours do not. The one source detected in the submillimetre waveband has narrow emission lines at a redshift of 1.45. Although the bulk of its radio emission originates from processes other than star formation, it is most probably a radio-quiet ultraluminous infrared galaxy.